

Application Advice

Winegrowing

Conventional agriculture As of May 2024



Planting out and spreading

Application rate for planting out

Water	100 l
AKRA Saat-Power	1 l

⚙️ Leave seedlings in the solution for 2–4 hours (after root pruning)

Spreading

AKRA Kombi	250 kg
AKRA DGC	As needed depending on fractional soil analyses

Application

Start of bud-swell phase

Water	200 l
AKRA Sulpur+	3 l
AKRA WD	2 l

Before flowering

Water	300 l
AKRA Plus 9	0.5 l
AKRA Blatt	2 l
Fungicide	When needed (50 % of the recommended application rate)
AKRA MSB	1 l
AKRA Azotobacter	0.5 l
AKRA Sulpur+	2 l
AKRA WD	0.2 l
Urea	10 kg

After flowering

Water	300 l
AKRA Plus 9	0.5 l
AKRA Blatt	2 l
Fungicide	When needed (50 % of the recommended application rate)
AKRA MSB	1 l
AKRA Azotobacter	0.5 l
AKRA Sulpur+	2 l
AKRA WD	0.2 l
Urea	10 kg

Spraying after 14 days or depending on infection pressure

Water	400 l
AKRA Plus 9	0.4 l
Fungicide	When needed (50 % of the recommended application rate)
AKRA MSB	2 l
AKRA Sulpur+	4 l
AKRA WD	0.4 l (increase by 0.4 % in the event of a spotted wing drosophila infestation)

INFO

- ⚙ Application up to 4 weeks max. before harvest.
- ⚙ The pH value of the spray solution should be adjusted to 5.5 with citric acid.
- ⚙ The addition of **AKRA** foliar fertiliser can and may reduce the pH value of the spray solution.
- ⚙ Mix in any other additives once the pH value has been adjusted.
- ⚙ Make sure that you add the additional products in the right order.
- ⚙ Do not spray in direct sunlight.
- ⚙ The dosage of **AKRA** foliar fertiliser must be adjusted to the quantity of water used.

SUCCESS

 WITH THE **AKRA FERTILISING SYSTEM**

» Guaranteed


 retrieve application
 advice online
